



How to help patients obtain MuGard® Oral Mucoadhesive

It is recommended that patients obtain *MuGard* prior to the start of their cancer treatment. *MuGard* Patient Reimbursement and Support Center works to deliver *MuGard* directly to patients.

To order *MuGard*, follow these two steps:

1

Prescribing physician completes and signs the Prescription Form in the Patient Starter Kit (prescription order form can also be downloaded from MuGard.com).

2

Completed form is faxed to
888-307-0071 or 929-977-0466

Let us know how much is needed; we'll handle the rest

MuGard is used 4 to 6 times a day throughout therapy. So one bottle lasts about a week. When completing the Prescription Form, the prescriber should indicate the initial quantity to be dispensed (**typically six 8-oz bottles**). Additional refills may be required depending on the specific regimen and length of treatment. If you have questions, call **toll-free at 866-900-5634**.



To learn more about the *MuGard* ordering process, see the other side of this sheet.

Important information for patients

Soleva Assist will contact patients directly by phone to verify their information. **It is important that patients speak with the Soleva Assist representative.** In the event that patients miss the call, **it is essential that they call Soleva Assist back as soon as possible at 866-900-5634**. Without this return call, timely dispatch of *MuGard* cannot be ensured.

Indication and Important Safety Information

Indication: MuGard® Oral Mucoadhesive is indicated for the management of oral mucositis/stomatitis (that may be caused by radiotherapy and/or chemotherapy) and all types of oral wounds (mouth sores and injuries), including aphthous ulcers/canker sores and traumatic ulcers, such as those caused by oral surgery or ill-fitting dentures or braces.

Contraindications: *MuGard* is contraindicated in patients with known hypersensitivity to any of the ingredients in the formulation.

Special Precautions for Use: Patients should avoid eating or drinking for at least one hour after using *MuGard*. After use, patients should replace the bottle cap and tightly seal the bottle. This product should not be used after the expiration date shown on the carton and product label. Do not use this product in patients with known sensitivity to any of the product's ingredients. Dilution of the product prior to use is not recommended.

Please see accompanying full Prescribing Information, also available at MuGard.com.


Working together to manage the suffering caused by oral mucositis





How Soleva Pharma, LLC. helps patients obtain MuGard® Oral Mucoadhesive

Soleva Assist Patient Reimbursement and Support Center (*Soleva Assist*)



Soleva Pharma, LLC. | Robbinsville, NJ 08691

Soleva Pharma and the logo designs presented in this material are trademarks of Soleva Pharma, LLC.

MuGard is a trademark of Soleva Pharma.

©2020 Soleva Pharma, LLC. DN-1255-0630